



Post-Operative for Wisdom Teeth Removal

Post-operative care is very important. Unnecessary pain and the complications of infection and swelling can be minimized if the instructions are followed carefully:

- Bite down on gauze until bleeding stops (about 1-2 hours)
- Swelling can be expected. Use ice packs on the surgical area (side of face) for the first 48 hours; apply ice 20 minutes on and 10 minutes off. (Note: Bags of frozen peas work well.) Keeping head elevated for 3 days. Swelling peaks on the second or third post-operative day.
- LIQUID DIET day of surgery (nothing with food particles) (i.e.- shakes, broth, yogurt, pudding, jello); soft food diet days after surgery (i.e..- Mashed potatoes, bread, pasta).
- Difficulty in opening your mouth widely and discomfort upon swallowing should be anticipated.
- Numbness of the lips and/or tongue on the affected side may be experienced for a variable period of time.
- If the corners of the mouth are stretched out, they may dry and crack. Your lips should be kept moist with cream or ointment.
- There will be a space where the tooth was removed. After day 5, Rinse your mouth with Peridex (twice a day) for 7 days after surgery. On day 5 after surgery, gently irrigate sockets with Peridex for 7 days after surgery. This cavity will gradually fill in with new tissue.
- Keep your fingers and tongue away from the socket or surgical area.
- For discomfort, take ibuprofen every 6 hours for 1 week.
- Drink plenty of fluids. (Do not use a straw—this creates suction in the mouth that could cause complications.)
- We suggest that you do not smoke for at least 5 days after surgery.



KC DENTAL IMPLANTS
& ORAL SURGERY

Dr. Brent Newby, DDS, MD | Dr. Brian Pannell, DDS | Dr. Taylor Markle, DDS | Dr. Lawson Parker, DMD |
Dr. Solon Kao, DDS | Dr. Matthew Wendahl, DMD, MD | Dr. Krista Lerner, DDS

- Avoid strenuous activity and do not exercise for at least 3–4 days after surgery.